

Photoinitiators as Reducing Agents

Georg Gescheidt¹, Hansjörg Grützmacher², Max Schmallegger¹

¹Institute of Physical and Theoretical Chemistry, TU Graz Stremayrgasse 9, A-8010 Graz,

²Laboratory of Inorganic Chemistry, ETH Zürich, Vladimir-Prelog-Weg 1-5, CH-8093 Zürich

In the recent time photoinitiators have found use as (chemically) photo-reducing agents.

In my contribution I will illustrate the “history” of these applications and report on mechanistic aspects and applications.

References

1. “Bis(acyl)phosphine Oxides as Stoichiometric Photo-Reductants for Copper Nanoparticle Synthesis: Efficiency and Kinetics”: M. Schmallegger, H. Grützmacher, G. Gescheidt, *Chemphotochem* **2022**, *6*. DOI: <https://doi.org/10.1002/cptc.202200155>
2. “Unprecedented Bifunctional Chemistry of Bis(acyl)phosphane Oxides in Aqueous and Alcoholic Media”: M. Schmallegger, A. Eibel, J. P. Menzel, A.-M. Kelterer, M. Zalibera, C. Barner-Kowollik, H. Gruetzmacher, G. Gescheidt, *Chem. Eur. J.* **2019**, *25*, 8982-8986. DOI: <https://doi.org/10.1002/chem.201900935>
3. “Benzil/triethylamine: a photo-reducing system for Cu²⁺”: M. Schmallegger, G. Gescheidt, *Monatsh. Chem.* **2018**, *149*, 499-504. DOI: <https://doi.org/10.1007/s00706-017-2085-7>
4. “The Redox Chemistry of Mono- and Bis(acyl)phosphane Oxides”: M. Zalibera, P.-N. Stebe, K. Dietliker, H. Grützmacher, M. Spichy, G. Gescheidt, *Eur. J. Org. Chem.* **2014**, *2014*, 331-337. DOI: <https://doi.org/10.1002/ejoc.201301214>